

UNCERTAINTY, TRAJECTORIES, AND DUALITY

ROBERT CARROLL
UNIVERSITY OF ILLINOIS, URBANA, IL 61801

To a wood nymph in an enchanted forest

ABSTRACT. First we show explicitly how uncertainty can arise in a trajectory representation. Then we show that the formal utilization of the WKB like hierarchy structure of dKdV in the description of (X, ψ) duality does not encounter norm constraints.

1. BACKGROUND

In a previous paper [4] (working with stationary states and ψ satisfying the Schrödinger equation (SE) **(A0)** $-(\hbar^2/2m)\psi'' + V\psi = E\psi$) we suggested that the notion of uncertainty in quantum mechanics (QM) can be phrased as incomplete information. The background theory here is taken to be the trajectory theory of Bertoldi-Faraggi-Matone-Floyd (cf. [1, 2, 3, 5, 11, 12, 13, 14]). The idea in [4] was simply that Floydian microstates satisfy a third order quantum stationary Hamilton-Jacobi equation (QSHJE)

$$(1.1) \quad \frac{1}{2m}(S'_0)^2 + \mathfrak{W}(q) + Q(q) = 0; \quad Q(q) = \frac{\hbar^2}{4m}\{S_0; q\};$$

$$\mathfrak{W}(q) = -\frac{\hbar^2}{4m}\{\exp(2iS_0/\hbar); q\} \sim V(q) - E$$

where **(A1)** $\{f; q\} = (f'''/f') - (3/2)(f''/f')^2$ is the Schwarzian and S_0 is the Hamilton principle function. If one recalls that the EP of Faraggi-Matone can only be implemented when $S_0 \neq \text{const}$ one may think of **(A2)** $\psi = R \exp(iS_0/\hbar)$ with $Q = -\hbar^2 R''/2mR$ and $(R^2 S'_0)' = 0$ where **(A3)** $S'_0 = p$ and $m_Q \dot{q} = p$ with $m_Q = m(1 - \partial_E Q)$ and $t \sim \partial_E S_0$. Thus microstates require three initial or boundary conditions in general to determine S_0 whereas the SE involves only two such conditions. Hence in dealing with the SE in the standard QM Hilbert space formulation one is not using complete information about the “particles” described by microstate trajectories. The price of underdetermination is then uncertainty in q, p, t for example. In the present note we will make this more precise and add further discussion.

2. SOME CALCULATIONS

It is shown in [12] that one has generally a formula

$$(2.1) \quad e^{2iS_0(\delta)/\hbar} = e^{i\alpha} \frac{w + i\bar{\ell}}{w - i\ell}$$

Date: August, 2003.
email: rcarroll@math.uiuc.edu.

$(\delta \sim (\alpha, \ell))$ with three integration constants, α, ℓ_1, ℓ_2 where $\ell = \ell_1 + i\ell_2$ and $w \sim \psi^D/\psi \in \mathbf{R}$. Note ψ and ψ^D are linearly independent solutions of the SE and one can arrange that $\psi^D/\psi \in \mathbf{R}$ in describing any situation. Here p is determined by the two constants in ℓ and has a form

$$(2.2) \quad p = \frac{\pm \hbar \Omega \ell_1}{|\psi^D - i\ell\psi|^2}$$

(where $w \sim \psi^D/\psi$ above and $\Omega = \psi'\psi^D - \psi(\psi^D)'$). Now let p be determined exactly with $p = p(q, E)$ via the Schrödinger equation and S'_0 . Then $\dot{q} = (\partial_E p)^{-1}$ is also exact so $\Delta q = (\partial_E p)^{-1}(\tau)\Delta t$ for some τ with $0 \leq \tau \leq t$ is exact (up to knowledge of τ). Thus given the wave function satisfying **(A0)**, with two boundary conditions at $q = 0$ say to fix uniqueness, one can create a probability density $|\psi|^2(q, E)$ and the function S'_0 . This determines p uniquely and hence $\dot{q}$. The additional constant needed for S_0 appears in (2.1) and we can write $S_0 = S_0(\alpha, q, E)$ since from (2.1) one has **(A4)** $S_0 - (\hbar/2)\alpha = -(i\hbar/2)\log(\beta)$ and $\beta = (w + i\bar{\ell})/(w - i\ell)$ with $w = \psi^D/\psi$ is to be considered as known via a determination of suitable ψ, ψ^D . Hence $\partial_\alpha S_0 = -\hbar/2$ and consequently **(A5)** $\Delta S_0 \sim \partial_\alpha S_0 \delta\alpha = -(\hbar/2)\Delta\alpha$ measures the indeterminacy in S_0 .

Let us expand upon this as follows. Note first that the determination of constants necessary to fix S_0 from the QSHJE is not usually the same as that involved in fixing $\ell, \bar{\ell}$ in (2.1). In particular differentiating **(A4)** in q one gets

$$(2.3) \quad S'_0 = -\frac{i\hbar\beta'}{\beta}; \quad \beta' = -\frac{2i\Re\ell w'}{(w - i\ell)^2}$$

Since $w' = -\Omega/\psi^2$ where $\Omega = \psi'\psi^D - \psi(\psi^D)'$ we get **(A6)** $\beta' = -2i\ell_1\Omega/(\psi^D - i\ell\psi)^2$ and consequently

$$(2.4) \quad S'_0 = -\frac{\hbar\ell_1\Omega}{|\psi^D - i\ell\psi|^2}$$

which agrees with p in (2.2) ($\pm\hbar$ simply indicates direction). We see that e.g. $S_0(x_0) = i\hbar\ell_1\Omega/|\psi^D(x_0) - i\ell\psi(x_0)|^2 = f(\ell_1, \ell_2, x_0)$ and $S''_0 = g(\ell_1, \ell_2, x_0)$ determine the relation between $(p(x_0), p'(x_0))$ and (ℓ_1, ℓ_2) but they are generally different numbers. In any case, taking α to be the arbitrary unknown constant in the determination of S_0 , we have $S_0 = S_0(q, E, \alpha)$ with **(A7)** $q = q(S_0, E, \alpha)$ and $t = t(S_0, E, \alpha) = \partial_E S_0$ (emergence of time from the wave function - ?). One can then write e.g. **(A8)** $\Delta q = (\partial q/\partial S_0)(\hat{S}_0, E, \alpha)\Delta S_0 = (1/p)(\hat{q}, E)\Delta S_0 = -(1/p)(\hat{q}, E)(\hbar/2)\Delta\alpha$ (for intermediate values $(\hat{S}_0, \hat{q})$) leading to

THEOREM 2.1. With p determined uniquely by two “initial” conditions so that Δp is determined and q given via (2.1) we have from **(A8)** the inequality **(A9)** $\Delta p \Delta q = O(\hbar)$ which resembles the Heisenberg uncertainty relation.

COROLLARY 2.1. Similarly **(A10)** $\Delta t = (\partial t/\partial S_0)(\hat{S}_0, E, \alpha)\Delta S_0$ for some intermediate value $\hat{S}_0$ and hence as before **(A11)** $\Delta E \Delta t = O(\hbar)$ (ΔE being precise).

3. EMBELLISHMENT FOR ENHANCED DKDV

In [2, 3, 5] we developed an enhanced dispersionless KdV theory to deal with the (X, ψ) duality of Faraggi-Matone [11] for the SE **(A0)**. The fact that connection of such diverse equations as SE and KdV could arise is simply based on the fact that they both share a second order differential equation. We want to indicate here more precisely this common feature and will summarize some material from [2, 3, 5, 6, 7, 8, 9] in the process.

3.1. (X, ψ) DUALITY. Consider a Schrödinger equation **(A12)** $[-(\hbar^2/2m)\partial_X^2 + V(X)]\psi = E\psi$ where E is in the spectrum (V and E are real and large X is used for reasons indicated below). Then let ψ^D be a second linearly independent solution of **(A12)** and define a pre-potential $\mathfrak{F}_E = \mathfrak{F}$ via **(A13)** $\psi^D = \partial\mathfrak{F}(\psi)/\partial\psi$ with **(A14)** $\partial_X\mathfrak{F} = \psi^D\partial_X\psi$ following [11, 12]. One expects **(A15)** $\mathfrak{F} = (1/2)\psi\psi^D + G$ with

$$(3.1) \quad \partial_X\mathfrak{F} = \psi^D\partial_X\psi = \frac{1}{2}(\partial_X\psi)\psi^D + (1/2)\psi\partial_X\psi^D + G_X \Rightarrow G_X = \frac{1}{2}(\psi^D\partial_X\psi - \psi\partial_X\psi^D) = \frac{1}{2}W$$

where the Wronskian W is constant. Hence take $G = (1/2)WX + c$ with $c = 0$ permitted to get **(A16)** $\mathfrak{F} = (1/2)\psi\psi^D + (1/2)WX$. We will consider situations where $\psi^D \sim \bar{\psi}$ with V and E real, and first, changing a minus sign in [2, 3, 5] to plus for a neater notation, we scale W via **(A17)** $W = \psi'\bar{\psi} - \psi\bar{\psi}' = 2i/\epsilon$ where $\epsilon \sim \hbar/\sqrt{2m}$. (cf. Remark 3.4 a description of scaling and normalization). Then, defining **(A18)** $\phi = \partial\mathfrak{F}/\partial\psi^2 = \bar{\psi}/2\psi$ one has a Legendre transform pair

$$(3.2) \quad -\frac{X}{i\epsilon} = \psi^2\partial_{\psi^2}\mathfrak{F} - \mathfrak{F}; \quad -\mathfrak{F} = \phi\frac{X_\phi}{i\epsilon} - \frac{X}{i\epsilon}$$

where $(X_\phi/i\epsilon) = \psi^2$ must be stipulated. Thus **(A20)** $\mathfrak{F} = (1/2)\psi\bar{\psi} - (X/i\epsilon)$, where it is interesting to note that $X/\epsilon = x$ in Section 3.1. This leads to **(A21)** $\mathfrak{F}_{\psi\psi\psi} = \frac{(E-V)}{4}(\mathfrak{F}_\psi - \psi\mathfrak{F}_{\psi\psi})^3$, which agrees with previous calculations in [2, 3, 5] when $\epsilon \rightarrow -\epsilon$.

REMARK 3.1. Equation **(A21)** can be written out in a different manner in terms of $\mathfrak{F}' = \partial\mathfrak{F}/\partial X$ in the form **(A22)** $\epsilon^2\mathfrak{F}''' - 2V'(\mathfrak{F} + \frac{X}{i\epsilon}) + 4(E-V)(\mathfrak{F} + \frac{1}{i\epsilon}) = 0$. This resembles a classical Gelfand-Dickey resolvent equation from soliton theory (cf. [10]). Indeed the resolvent equation arises in the form **(A23)** $R''' + 4uR' + 2u'R + 4z^2R' = 0$ in reference to a Lax operator $L = \partial^2 + u$ and R is the formal analogue of the diagonal of the kernel of a resolvent (i.e. an operator inverse to $\partial^2 + u + z^2$ corresponding to a Green's function. Such an operator can be constructed as the product of two solutions of $(\partial^2 + u + z^2)\psi = 0$ (i.e. $R = \psi_1\psi_2$ which in the present situation refers to $R = \psi\bar{\psi}$). Thus $\psi\bar{\psi} = \Xi \sim 2\mathfrak{F} + (2X/i\epsilon) = 2(\mathfrak{F} + (X/i\epsilon))$ with **(A12)** $\sim$ **(A12')** $\psi'' - (1/\epsilon^2)V\psi + (1/\epsilon^2)E\psi = 0$ so $u \sim -(V/\epsilon^2)$ and $z^2 \sim E/\epsilon^2$ with $\mathfrak{F}''' = \Xi'''$. Hence **(A22)** amounts to $\Xi \sim R$ with

$$(3.3) \quad \Xi''' - \frac{2V'}{\epsilon^2}\Xi + \frac{4(E-V)}{\epsilon^2}\Xi' = 0 \equiv \Xi''' + 2u'\Xi + 4z^2\Xi' + 4u\Xi' = 0$$

This gives us a first direct connection between (X, ψ) duality and the KdV theory based on the fact that a very special second order differential equation **(A12)** $\sim$ **(A12')** appears in both situations ($\epsilon^2 \sim \hbar^2/2m$ - see below). Evidently such connections based on properties of an operator $\partial^2 + u$ are perfectly natural and mathematically well founded; in particular

we will use techniques from the theory of **(A12')** to describe quantities arising via **(A12)** in (X, ψ) duality. The fact that some of the techniques arise from KdV (or dKdV) theory via tau functions, Lax operators, etc. is purely incidental (and fortuitous). ■

3.2. DISPERSIONLESS THEORY FOR KP and KDV. We assume the standard KP and KdV theory is known (cf. [5, 6, 10]). Then one can think of fast and slow variables with $\epsilon x = X$ and $\epsilon t_n = T_n$ so that $\partial_n \rightarrow \epsilon \partial / \partial T_n$ and $u(x, t_n) \rightarrow U(X, T_n)$ to obtain from the KP equation $(1/4)u_{xxx} + 3uu_x + (3/4)\partial^{-1}\partial_x^2 u = 0$ the equation $\partial_T U = 3U\partial_X U + (3/4)\partial^{-1}(\partial^2 U / \partial T^2)$ when $\epsilon \rightarrow 0$ ($\partial^{-1} \rightarrow (1/\epsilon)\partial^{-1}$). In terms of hierarchies the theory can be built around the Lax operator $L = \partial + \sum_{n=1}^{\infty} u_{n+1}\partial^{-n}$. Then writing (t_n) for (x, t_n) (i.e. $x \sim t_1$ here) consider **(A23)** $L_\epsilon = \epsilon \partial + \sum_{n=1}^{\infty} u_{n+1}(\epsilon, T)(\epsilon \partial)^{-n}$. Now one assumes $u_n(\epsilon, T) = U_n(T) + O(\epsilon)$, etc. and sets (recall $L\psi = \lambda\psi$)

$$(3.4) \quad \psi = \left[1 + O\left(\frac{1}{\lambda}\right)\right] \exp\left(\sum_1^{\infty} \frac{T_n}{\epsilon} \lambda^n\right) = \exp\left(\frac{1}{\epsilon} S(T, \lambda) + O(1)\right);$$

where **(A24)** $\tau = \exp(1/\epsilon^2 F(T) + O(1/\epsilon))$. Note for the approximation of potentials one assumes e.g. $v = v(x, t_i) \rightarrow v(X/\epsilon, T_i/\epsilon) = V(X, T_i) + O(\epsilon)$. This is standard in dispersionless KP = dKP and certainly realized by quotients of homogeneous polynomials for example. In fact it is hardly a restriction since we are primarily interested here in the X dependence and given e.g. $F(X) = \sum_0^{\infty} a_n X^n$ consider $\tilde{F}(x, t_i) = a_0 + \sum_1^{\infty} (x^n / \prod_2^{n+1} t_i)$. Then $\tilde{F}(X/\epsilon, T_i/\epsilon) = a_0 + \sum_1^{\infty} (X^n / \prod_2^{n+1} T_i)$ and one can choose the T_i recursively so that $1/T_1 = a_1$, $1/T_1 T_2 = a_2, \dots$, leading to $F(X) = \tilde{F}(X, T_i)$. Thus one can work with a $t \sim T$ dependent theory and eventually capture the original $F(X)$ for a stationary theory by specializing and freezing the T_i .

REMARK 3.2. We recall also that $\partial_n L = [B_n, L]$, $B_n = L_+^n$, $L\psi = \lambda\psi$, and in terms of tau functions $\psi = \tau(T - (1/n\lambda^n)) \exp[\sum_1^{\infty} T_n \lambda^n] / \tau(T)$. Putting in the ϵ and using ∂_n for $\partial / \partial T_n$ now, with $P = S_X$, one obtains **(A25)** $\lambda = P + \sum_1^{\infty} U_{n+1} P^{-n}$ and $P = \lambda - \sum_1^{\infty} W_i \lambda^{-1}$ (when $\epsilon \rightarrow 0$) There are then many beautiful formulas for dKP (see e.g. [5]) but we emphasize here that in contrast to dKdV we are not going to let $\epsilon \rightarrow 0$; instead we use it to balance terms below. ■

Now look at the dispersionless theory for KdV based on k where $\lambda^2 \sim (\pm ik)^2 = -k^2$. When $\epsilon \rightarrow 0$ one obtains for $P = S_X$, $P^2 + q = -k^2$, and we write $\mathcal{P} = (1/2)P^2 + p = (1/2)(ik)^2$ with $q \sim 2p \sim 2u_2$. One has $\partial k / \partial T_{2n} = \{(ik)^{2n}, k\} = 0$ and from $ik = P(1 + qP^{-2})^{1/2}$ we obtain

$$(3.5) \quad ik = P \left(1 + \sum_1^{\infty} \binom{\frac{1}{2}}{m} q^m P^{-2m}\right)$$

($\epsilon \rightarrow 0$ here and we use $+ik$ since it is appropriate later). We refer now to [5, 7, 8, 15] for dispersionless KP (= dKP) and consider here $\psi = \exp[(1/\epsilon)S(X, T, \lambda)]$ (note **(A26)** $L^2 = \partial_x^2 - v \rightarrow \epsilon^2 \partial_X^2 - V$). Thus $P = S' = S_X$ with $P^2 = V - E$ (when $\epsilon \rightarrow 0$) and $E = \pm \lambda^2$ real will involve us in a KdV situation and some routine calculation yields; recall $X_\psi = 1/\psi'$

with $\psi' = (P/\epsilon)\psi$ (cf. [5, 8]). This leads to

$$(3.6) \quad \Im \mathfrak{F} = \frac{X}{\epsilon}; \quad \Re \mathfrak{F} = \frac{1}{2}|\psi|^2 = \frac{1}{2}e^{\frac{2}{\epsilon}\Re S} = -\frac{1}{2\Im P}$$

In the present situation $|\psi|^2 = \exp[(2/\epsilon)\Re S]$ and $2\phi = \exp[-(2i/\epsilon)\Im S]$ can play the roles of independent variables and we see that (A27) $|\psi|^2 \Im P = 1$ while $\psi^2 \phi = (1/2)|\psi|^2$. Now note that for $L = \partial + \sum_1^\infty u_i \partial^{-i}$, $L_+^2 = \partial^2 + 2u_1$, and $u_1 = \partial^2 \log(\tau)$ where τ is the famous tau function. From (A26) this implies $v = -2\partial^2 \log(\tau)$ here, from which $V = -2F_{XX}$ for $\tau = \exp[(1/\epsilon^2)F + O(1/\epsilon)]$ in the dispersionless theory (cf. [5, 8]). Then writing out the Gelfand-Dickey resolvent equation (A22) yields

$$(3.7) \quad \epsilon^2 \mathfrak{F}''' + \left(\mathfrak{F}' + \frac{1}{i\epsilon} \right) (8F'' + 4E) + 4F''' \left(\mathfrak{F} + \frac{X}{i\epsilon} \right) = 0$$

which provides a relation between F and $\mathfrak{F}$. This is interesting since F plays the role of a prepotential or free energy in the dKdV theory (cf. [2, 3, 5]). We recall below how to embellish all this with an ϵ -modification (or enhancement) of dKP and dKdV. Note that there is no neglect of $O(\epsilon)$ terms in (3.7) and $X/i\epsilon = x$ should be well defined which suggests a scale dependence in QM and perhaps the emergence of space from the wave function.

Next one has $\Im \mathfrak{F} = X/\epsilon$ and $|\psi|^2 = \exp[(2/\epsilon)\Re S]$. In order to have $|\psi|^2 \leq 1$ as a fundamental variable in the (X, ψ) theory some control over $\Re S$ is needed and this problem is resolved below in Remark 3.5 and Proposition 3.1. One notes that dKdV (with $\epsilon \rightarrow 0$ here) involves $\lambda = \pm ik = P(1 + \sum_1^\infty U_m P^{-2m})$ with $\lambda^2 = -k^2$ real for k real (as in (3.6)). In order to satisfy $|\psi|^2 \Im P = 1$ we will want $\lambda = ik$ so that $Q = \Im P$ is positive. The U_m are real so $P = iQ$ corresponds to k real and $(ik)_+^{2n+1}$ will be purely imaginary (only “times” t_{2n+1} arise in KdV). In addition $\Re S = \Re[\sum_0^\infty T_{2n+1}(ik)^{2n+1} + \sum_1^\infty S_{j+1}(ik)^{-j}] = 0$ for k real since $S_{j+1} = -\partial_j F/j$ with $\partial_{2n} F = 0$. Now this would imply $|\psi|^2 = 1$, which is absurd, so we introduce an expansion $S \rightarrow \tilde{S} = \sum \epsilon^j S^j$ (in particular this takes into account the $O(1)$ terms in $\psi = \exp[(1/\epsilon)S + O(1)]$). Note that such $O(1)$ terms (and others) arise quite naturally from the vertex operator equation (VOE) via $\log \psi = (S/\epsilon) + O(1) = \log \tau[\epsilon, T_n - (\epsilon/n\lambda^n)] - \log \tau + \sum_1^\infty T_n \lambda^n / \epsilon$ with $\log \tau = (F/\epsilon^2) + O(1/\epsilon)$ and $S_{n+1}^0 = -(\partial_n F/n)$. Thus e.g.

$$(3.8) \quad F\left(T_n - \frac{\epsilon}{n\lambda^n}\right) - F(T_n) = -\epsilon \sum_1^\infty \left(\frac{\partial_n F}{n\lambda^n}\right) + \frac{\epsilon^2}{2} \sum \left(\frac{F_{mn}}{mn}\right) \lambda^{-m-n} + O(\epsilon^3)$$

This leads to first terms of the form $\tilde{S} = S^0 + \epsilon S^1$ with

$$(3.9) \quad S^1 = \frac{1}{2} \sum \left(\frac{F_{mn}}{nm}\right) \lambda^{-m-n}; \quad S_X \sim P + \frac{\epsilon}{2} \sum \left(\frac{F_{1mn}}{nm}\right) \lambda^{-m-n}$$

Eventually we will want also an embedding $F \rightarrow \tilde{F} = \sum \epsilon^j F^j$ as well. With this development one finds as a first approximation in ϵ

$$(3.10) \quad |\psi|^2 = e^{2\Re S^1} = \exp \left[\Re \sum \left(\frac{F_{mn}^0}{mn}\right) \lambda^{-m-n} \right]$$

where only terms $F_{2m+1,2n+1}^0 \lambda^{-2(m+n)-2}$ arise and these will be real for k real (so $|\psi|^2 \leq 1$ becomes tenable). We refer to Remark 3.4 and Proposition 3.1 where it is shown that this can always be achieved without constraints. Thus (A28) S^0 and $P^0 = S_X^0 = iQ$ are imaginary while S^1 and $P^1 = S_X^1$ are real.

REMARK 3.3. We note from [12] that a standard WKB approach to $(\hbar^2/2m)\psi'' + (E - V)\psi = 0$ involves $\psi = A \exp(is/\hbar)$ where s and A are even functions of ϵ . This leads to (A29) $(s')^2 - 2m(E - V) = \hbar^2 \frac{A''}{A}$; $2A's' + As'' = 0$. Thus (A30) $A = c(s')^{-1/2}$ and the first equation becomes (A31) $(s')^2 = 2m(E - V) - \frac{\hbar^2}{2} \{s; x\}$; $\{s; x\} = \frac{s'''}{s'} - \left(\frac{s''}{s'}\right)^2$. We recall that the (X, ψ) duality theme was based on $-\epsilon^2 \psi'' + V(X)\psi = E\psi$ with $\mathfrak{F} = (1/2)\psi\bar{\psi} - (X/i\epsilon)$ and $\psi = \exp[\tilde{S}/\epsilon]$ was employed ($\epsilon = \hbar/\sqrt{2m}$) with (A32) $\tilde{S} = \sum_0^\infty \epsilon^j S^j$ (S^{2j+1} real and S^{2j} imaginary for k real, $\lambda = ik$). This is related (in an expanded dKdV theory) to $\tilde{s} \sim \sqrt{2m}\mathfrak{F}\tilde{S}$ and $\log A = (1/\epsilon)\Re\tilde{S}$. Also since ψ and $\bar{\psi}$ satisfy the Schrödinger equation we know that the square eigenfunctions $\psi\bar{\psi}$, ψ^2 , and $(\bar{\psi})^2$ satisfy the GD equation. Further from [12] a general solution of (A31) is given by $s' = \pm\sqrt{2m}(a\psi^2 + b\bar{\psi}^2 + c\psi\bar{\psi})^{-1}$ where ψ , $\bar{\psi}$ are normalized solutions of (A32) so we introduce a constraint (A33) $\int |\psi|^2 dX = 1$. Here one recalls that $|\psi|^2 = 1/\mathfrak{F}\tilde{P} = \exp[(1/\epsilon)\Re\tilde{S}] = \exp[\sum_0^\infty \epsilon^{2j} S^{2j+1}]$ and evidently one will have to scale $\psi \rightarrow c\psi$ in order to insure (A33); this amounts to a normalization of a BA function and we will show that this introduces no problems (cf. Remark 3.5). Now the GD equation (3.3) for $\mathfrak{F}$ can be written in terms of $\Xi = \psi\bar{\psi}$, or ψ^2 , or $\bar{\psi}^2$ as (A34) $\epsilon^2 \Xi''' - 4V\Xi' - 2V'\Xi + 4E\Xi' = 0$ so given s' as above one sees the connection via $s' = c/|\psi|^2$ satisfies (A31) while $|\psi|^2 = c/s'$ satisfies (A34) (for any c). We have also shown incidentally that $|\psi|^2 s' = c$ is a consequence of the second order equation (A12) $\sim$ (A12') and thus arises from the WKB approach (independently of (X, ψ) duality). ■

In [2, 3, 5] we employed a full connection to dKdV by using the tau function (A35) $\tau = \exp[(1/\epsilon^2)F(T) + O(1/\epsilon)]$ with (A36) $V = -2F_{XX} = -2F''$. Then the GD equation takes the form (A37) $\epsilon^2 \Xi''' + 4F''' \Xi + 4E\Xi' + 8F'' \Xi' = 0$. It is in fact F , with its fundamental coefficients (A38) $F_{mn} = \partial_m \partial_n F = \partial^2 F / \partial T_m \partial T_n$ and their relations to Hirota theory, which make the enhanced dKdV theory interesting and useful. We recall here the expansions (A39) $\tilde{S} = \sum \epsilon^j S_j$, $\tilde{F} = \sum \epsilon^j F_j$, $\tilde{P} = \tilde{S}_X = \sum \epsilon^j P_j$ which from the SE $\epsilon^2 \psi'' + 2F''\psi = -E\psi$ and $\epsilon^2 \psi_{XX} = \epsilon \tilde{P}_X \psi + \tilde{P}^2 \psi$ yield

$$(3.11) \quad \left(\sum_0^\infty \epsilon^j P_j \right)^2 + \epsilon \sum_0^\infty \epsilon^j P_j' + 2 \sum_0^\infty \epsilon^j F_j'' = -E = -k^2$$

(we use sub or superscripts according to notational convenience and will write S^0 in order to distinguish this from our original $S_0 \sim s$). This leads to (A40) $P_0^2 + 2F_0'' = -E$, $2P_0 P_1 + P_0' + 2F_1'' = 0$, $P_1^2 + 2P_0 P_2 + P_1' + 2F_2'' = 0, \dots$ (there is also another (equivalent) way to determine coefficients indicated in [2, 3, 5]). Take now $F_{2j+1} = 0$ (for consistency of equations - cf. [2, 3, 5]); this only stipulates what kind of ϵ extensions of F are compatible with (X, ψ) duality in an extended sense, namely (A41) $\exp[(2/\epsilon)\Re\tilde{S}]\mathfrak{F}\tilde{P} = 1$ (which also appears as an essential ingredient for using WKB type methods). Then from (3.11) one obtains (A42) $P_k^2 + 2 \sum_0^{k-1} P_i P_{2k-i} + P_{2k-1}' + 2F_{2k}'' = 0$; $2 \sum_0^k P_i P_{2k+1-i} + P_{2k}' = 0$ (cf.

[2, 3, 5]). In particular (A43) $P_0^2 + 2F_0'' = -E$; $2P_0P_1 + P_0' = 0$; $P_1^2 + 2P_0P_2 + P_1' + 2F_2'' = 0$ with $P_2' + 2(P_0P_3 + P_1P_2) = 0; \dots$. Consequently (A44) $P_1 = -P_0'/2P_0$ and one can recover all P_{2j} from E and F_{2m}'' . Hence s' is determined by E and the F_{2m}'' and this will be a solution of an enhanced QSHJE. In terms of construction we note that (A45) $P_0^2 = -2F_0'' - E \sim V - E$ and via (A44) one has $2P_0'P_1 + 2P_0P_1' + P_0'' = 0$ so (A46) $2P_0P_2 = -2F_2'' - P_1' - P_1^2 = -2F_2'' + \frac{1}{2} \left[\left(\frac{P_0''}{P_0} \right) - \frac{3}{2} \left(\frac{P_0'}{P_0} \right)^2 \right]$. (note the Schwarzian representation $(1/2)\{S_0; q\}$ in (A46)). In any event V , E , and F_2'' determine P_2 . Continuing one can write (A47) $P_{2k} = f(V, E, F_{2m}'')$ ($1 \leq m \leq k$) and there are no integration constants involved. We note that even if $F_2'' = 0$ there is a correction term (A48) $P_2 = \{S_0; q\}/4P_0$.

REMARK 3.4. Let us examine again (A49) $|\psi|^2 \leq 1$ and (A57) $|\psi|^2 \Im P = 1$. First (A50) is fundamental to (X, ψ) duality and in fact to the whole idea of a WKB type expression $\psi = \exp(S/\epsilon)$. Thus write the solution of (A51) $(\epsilon^2 \partial_X^2 - V)\psi = -k^2 \psi$ as $\psi = R \exp(is/\epsilon)$ so that (A52) $\psi' \bar{\psi} = R'R + iR^2 s'/\epsilon$. Also (A51) implies (A53) $(s')^2 + V - E - (\epsilon^2 R''/R) = 0$ with $\partial(R^2 s') = 0$. In particular (A54) $R^2 s' = c$ (constant) and since $s' \sim \Im S_X = \Im P$ (or $\Im \tilde{P}$) we have $|\psi|^2 \Im \tilde{P} = c$ by virtue of the WKB formulation. Note also from $(\epsilon^2 \partial_X^2 - V)\bar{\psi} = -k^2 \bar{\psi}$ one has for $\bar{\psi} = R \exp(-is'/\epsilon)$ the formula $\bar{\psi}' \psi = R'R - (is'R^2/\epsilon)$. Combining this with (A52) leads to (A55) $W = \psi' \bar{\psi} - \bar{\psi}' \psi = 2iR^2 s'/\epsilon = (2i/\epsilon)c$. Thus the normalization of W involves an integration constant c so one could say that c is determined by the normalization of W . Note however that using $R^2 s' = c$ in rewriting the “quantum potential” $Q = -\epsilon^2 R''/R$ in terms of s' we have $2RR's' + R^2 s'' = 0$ with (A56) $\frac{R''}{R} = -\frac{1}{2} \left[\frac{s'''}{s'} + \frac{3}{2} \left(\frac{s''}{s'} \right)^2 \right]$. Hence (A57) $Q = (\epsilon^2/2)\{s; X\}$ and the constant disappears in constructing the QSHJE - which then requires three “initial” conditions for integration. The important point here is that (A58) $|\psi|^2 = R^2 = c/s'$ with c a priori an integration constant and in addition c scales the Wronskian as in (A55). We could also simply scale ψ via $\psi \rightarrow \sqrt{\omega} \psi$ to achieve the same result for $\omega \sim c$. This is interesting since c disappears in the QSHJE and loses its status of integration constant. Therefore we can think of c as a scaling factor ($c \sim \omega$) in the projective ray representation of ψ . This does not however fix the theory to take place on a sphere of radius $|\psi|^2 = \omega$ but appears to be a harmless scaling which can in fact be used to force $|\psi|^2 \leq 1$ and we will have (A54) to measure this. In any case (A59) $|\psi|^2 = \exp[2\Re \tilde{S}/\epsilon] = \exp[2 \sum \epsilon^{2j} P_{2j+1}]$ and we are free to scale ψ so for $\psi \rightarrow \sqrt{\omega} \psi$ with $|\psi|^2 \rightarrow \omega |\psi|^2$ the requirement is (A60) $\log(\omega |\psi|^2) = \log |\psi|^2 + \log \omega = \log \omega + 2 \sum_0^\infty \epsilon^{2j} P_{2j+1} \leq 0$. But for $\omega < 1$ one has $\log \omega < 0$ so given any magnitude for the series $2 \sum_0^\infty \epsilon^{2j} P_{2j+1}$ we can find ω such that $(\omega |\psi|^2) \leq 1$. ■

PROPOSITION 3.1. The apparent integration constant c in $R^2 s' = c$ can be thought of in terms of an arbitrary scaling constant $c = \omega$ for ψ and hence there need be no concern about satisfying $|\psi|^2 \leq 1$ in the enhanced dKdV theory.

REFERENCES

- [1] G. Bertoldi, A. Faraggi, and M. Matone, hep-th 9909201
- [2] R. Carroll, Nucl. Phys. B, 502 (1997), 561-593
- [3] R. Carroll, Lect. Notes Physics 502, Springer, 1998, pp. 33-56

- [4] R. Carroll, *Canad. Jour. Phys.*, 77 (1999), 319-325
- [5] R. Carroll, *Quantum theory, deformation, and integrability*, North-Holland, 2000
- [6] R. Carroll, *Topics in soliton theory*, North-Holland, 1991
- [7] R. Carroll, *Jour. Nonlin. Sci.*, 4 (1994), 519-544; *Teor. Mat. Fiz.*, 99 (1994), 220-225
- [8] R. Carroll and Y. Kodama, *Jour. Phys. A*, 28 (1995), 6373-6387
- [9] R. Carroll, *Proc. NEEDS '94*, World Scientific, pp. 29-33
- [10] L. Dickey, *Soliton equations and Hamiltonian structure*, World Scientific, 1991
- [11] A. Faraggi and M. Matone, *Phys. Rev. Lett.*, 78 (1997), 163-166
- [12] A. Faraggi and M. Matone, *Inter. Jour. Mod. Phys. A*, 15 (2000), 1869-2017
- [13] E. Floyd, *Inter. Jour. Mod. Phys. A*, 14 (1999), 1111-1124; 15 (2000), 1363-1378; *Found. Phys. Lett.*, 13 (2000), 235-251; quant-ph 0009070, 0302128 and 0307090
- [14] E. Floyd, *Phys. Rev. D*, 29 (1984) 1842-1844; 26 (1982), 1339-1347; 34 (1986), 3246-3249; 25 (1982), 1547-1551; *Jour. Math. Phys.*, 20 (1979), 83-85; 17 (1976), 880-884; *Phys. Lett. A*, 214 (1996), 259-265; *Inter. Jour. Theor. Phys.*, 27 (1998), 273-281
- [15] K. Takasaki and T. Takebe, *Rev. Math. Phys.*, 7 (1995), 743-808; *Inter. Jour. Mod. Phys. A*, Supplement 1992, pp. 889-922